

Group 2

Indicators Prioritization Matrix For The ASEAN Nutrition Surveillance System (ANSS)

ASEAN Member State in GROUP 2

ASEAN Member State	Name of Delegates
Brunei Darussalam	1. Mr. Zakaria Haji Kamis
	2. Mr. Abdul Al Bari
Cambodia	3. Dr. Chea Mary
Indonesia	4. Mrs. Andri Mursita
Lao PDR	5. Mr.Somchay Sithpangna
Malaysia	6. Mrs. Junidah Binti Raib
Philippines	7. Ms. Hygela Ceres Catalina B Gawe
Thailand	8 Dr Napaphan Viriyautsahakul, MD
	9. Dr Sunisa S, MD
ASEAN Secretariat	10. Alautiah (Tia)
Facilitator (Thailand)	11. Dr Saiyed M Hyder
Rapporteur 1	12. Dr Wannachanok Boonchoo (Bim)
Rapporteur 2	13. Ms Punnasa

Nutrition Targets	Malaysia	Brunei	Indonesia	Lao PDR	Cambodia	Philippines	Thailand
<p>Target 6</p> <p>(7) Proportion of infants 0-5 months of age who are fed exclusively with breast milk</p>	<p>✓</p> <p>Source: NHMS(2016)</p>	<p>✓✓</p> <p>Source: NHNSS and Bruhim MCH</p>	<p>✓</p> <p>Source: Basic Health Research survey (2007,2013,2018)</p>	<p>✓</p>	<p>✓</p> <p>Source: CDHS</p>	<p>✓</p> <p>Source: NNS</p>	<p>✓✓</p> <p>Source: MICS,HDC</p>
<p>Target7</p> <p>(8) Prevalence of wasting (Low weight-for-height) in children 0-59 months of age</p>	<p>✓</p> <p>Source: NHMS(2016)</p>	<p>✓✓</p> <p>Source: NHNSS and Bruhim MCH</p>	<p>✓</p> <p>Source: Basic Health Research survey (2007,2013,2018) Nutritional status survey in children 0 – 59 mo (annual)</p>	<p>✓</p>	<p>✓</p> <p>Source: CDHS</p>	<p>✓✓</p> <p>Source: NNS</p>	<p>✓✓</p> <p>Source: MICS,HDC</p>
<p>Target8</p> <p>(9) Prevalence of overweight (BMI≥25) and obesity (BMI≥30) in women 18+ years</p>	<p>✓</p> <p>Source: NHMS(2015)</p>	<p>✓</p> <p>Source: NHNSS (2010), STEPS (2016)</p>	<p>✓</p> <p>Source: Basic Health Research survey (2007,2013,2018) (obesity BMI ≥ 27)</p>	<p>✓</p>	<p>N/A</p>	<p>✓✓</p> <p>Source: NNS</p>	<p>✓✓</p> <p>Source: NHES,HDC</p>

Nutrition Targets	Malaysia	Brunei	Indonesia	Lao PDR	Cambodia	Philippines	Thailand
(10) Prevalence of overweight (BMI≥25) and obesity (BMI≥30) in men 18+ years	✓ Source: NHMS(2015)	✓ Source: NHNSS (2010), STEPS (2016)	✓ Source: Basic Health Research survey (2007,2013,2018) (obesity BMI ≥ 27)	✓	N/A	✓✓ Source: NNS	✓✓ Source: NHES,HDC
(11) Prevalence of overweight (>1SD body mass index for age and sex) and obesity (>2SD body mass index for age and sex) in school age children and adolescents 5-18 years of age	✓ Source: NHMS (2017), GBHS (2012)	✓ Source: NHNS(2010), GSHS(2014)	✓ Source: Basic Health Research survey (2007,2013,2018) (Age group: 5-12 and 13-15, and 16 - 18 y o, available by sex)	✓	N/A	✓✓ Source: NNS	✓✓ Source: NHES
(12) Age-standardized prevalence of raised blood glucose/diabetes among persons aged 18+	✓ Source: NHMS (2015)	✓ Source: STEPS (2016)	✓ Source: Routine data from facility based	✓	N/A	✓✓ Source: NNS	✓✓ Source: NHES, Bureau of NCD's Annual Report

Nutrition Targets	Malaysia	Brunei	Indonesia	Lao PDR	Cambodia	Philippines	Thailand
<p>Target9</p> <p>(13) Prevalence of insufficiently physically active adolescents (10-19 years old), defined as less than 60 minutes of moderate to vigorous intensity activity daily</p>	<p>✓</p> <p>Source: NHMS (2017)</p>	<p>✓</p> <p>Source: GSHS (2014)</p>	<p>Basic Health Research survey (2007,2013,2018)</p> <p>Age group 10 -14 and 15-19 y o cut off point physically active more than 150 minutes per week</p> <p>available by sex</p>	N/A	N/A	N/A	N/A
<p>(14) Age-standardized prevalence of insufficiently physically active person aged 18+ (defined as less than 150 minutes of moderate-intensity activity per week, or equivalent)</p>	<p>✓</p> <p>Source: NHMS (2015)</p>	<p>✓</p> <p>Source: GSHS (2014)</p>	<p>Basic Health Research (2007,2013,2018)</p> <p>Age group 20 – 24, 25 – 29, 30 – 34, 35 -39, 40-44, 45 – 49, 50-54, 55-59, 60 – 64, and 65+ y o</p> <p>available by sex</p>	N/A	N/A	<p>✓</p> <p>Source: NNS</p>	N/A
<p>Target10</p> <p>(15) Age-standardized mean population intake of salt (sodium chloride) per day grams in persons aged 18+ years</p>	<p>✓</p> <p>Source: NYCOSS (2017)</p>	<p>✓</p> <p>Source: STEPS (2016)</p>	<p>✓</p> <p>Source: Total Diet Survey (2014)</p> <p>age group 5-12, 13-18 and 19-55 yo, by place, by sex</p>	N/A	N/A	<p>✓</p> <p>Source: NNS</p>	N/A

Breakdown of sources

1. NHMS = National Health Nutrition Status
2. GBHS = Global Health Survey
3. NHNSS = National Health and Mobility Survey
4. CDHS = Cambodia Demographic and Health Survey
5. MICS = Multiple Indicators Cluster Survey
6. HDC = Health Data Center
7. NNS = National Nutrition Survey

Site Visit Summary (Cambodia)

Health Promotion Center 5, Ratchaburi Province: IDD Prevention and Control Programme:

Strength: Triferdine supplementation to all pregnant women from ANC 1 to 6 month after delivery in combination with the consumption of iodized salt really see the adequate iodine intake

Weakness: 100 % distribution rate, but not yet monitor the compliance. We do not see the result of the reduction of anemia, NTD and children intelligent (IQ)